

Developing an understanding of the risk factors for falls among adults with intellectual disabilities: a commentary

Commentary on: Pope J, Truesdale M, Brown M. Risk factors for falls among adults with intellectual disabilities: A narrative review. *J Appl Res Intellect Disabil.* 2021; 34: 274–285. <https://doi.org/10.1111/jar.12805>

Alison J Doherty,¹ Annette O’Donoghue² and Joanna Harrison¹

¹ Synthesis, Economic Evaluation and Decision Science (SEEDS) Group, Health Technology Assessment Unit, Applied Health Research Hub, University of Central Lancashire

² University Hospitals of Morecambe Bay NHS Foundation Trust

adoherty7@uclan.ac.uk

Falls are a common and serious problem for older people in the general population and the human costs include pain, loss of confidence, loss of independence and mortality (National Institute for Health & Care Excellence (NICE), 2013). Even if the person is not injured, they may have a fear of falling, which may result in avoidance of social activities, reduced mobility, and increased frailty (Public Health England (PHE), 2019). Falls also affect the carers of older people who fall (Ang et al, 2020), including costs in terms of carer time and absence from work (Snooks et al, 2011). Falls in the general population cost the National Health Service (NHS) more than £2.3 billion per year (NICE, 2013).

While there is evidence for older people at risk of falls in the general population, there is limited research investigating falls among those with intellectual disabilities (ID) who may share many of the same risk factors, and which may increase their risk of falls from a younger age (Cox et al, 2010). The

term ID (used throughout this paper) is an internationally recognised term to define people who may be classed as having learning disabilities (LD). ID refers to individuals with a significantly reduced ability to understand new or complex information and to learn and apply new skills (impaired intelligence), which started before adulthood and results in a reduced ability to cope independently (impaired social functioning) (World Health Organization (WHO), 2010). People with mild-to-moderate ID often live independently and are usually able to communicate most of their needs, but they may need support to understand complex issues. People with severe-to-profound ID may require a higher level of support. There are many conditions associated with ID, for example, Down’s syndrome, Williams syndrome, and Fragile X syndrome (Mencap, 2022). It is estimated that 1.5 million people in the UK have an ID (approximately 2% of the UK’s population), with the majority living independently in their local communities or with families or paid support (WHO, 2010; Mencap, 2022). However, the exact numbers are unknown, as they may not be registered with health or social care related services. Falls and fall-related injuries are a common and serious problem for people with ID. Research from various studies suggest that 25–40% of people with ID experience at least one fall per year (Hsieh et al, 2012; Petropoulou et al, 2017; Finlayson et al, 2010). Around one-third of falls involving people with intellectual disabilities result in injury, with the rate of fractures higher than in the general population, which may be due to an increased risk of osteoporosis (PHE, 2019). It is commonly understood that people with intellectual disabilities may be considered ‘older adults’ (from the age of 40 years onwards) (Hermans and Evenhuis, 2014). Given this, their risk of falls and subsequent

Abstract

Falls are a serious and common problem for people with intellectual disabilities (ID). Whilst there is available evidence on falls risk factors for the general population, there is a lack of awareness and understanding of the contributing risk factors for this population. This commentary critically appraises a recent narrative review which sought to identify the risk factors for falls among people with ID. Clinical practice implications: Community nurses may identify people with ID at risk of falls, and work alongside other healthcare professionals and carers to provide tailored multidisciplinary falls-prevention interventions for individuals with ID living in the community.

Keywords: Falls • intellectual disabilities • community nursing • risk factors

injury may occur earlier than in the general population and over a longer period of time (Cox et al, 2010; Pal et al, 2013; Oppewal et al, 2014).

Willgoss et al (2010) identified that risk factors for falls in people with ID may include older age, impaired mobility, epilepsy and behavioural problems. Evidence concerning risk factors for falls in older people within the general population may also be relevant to people with intellectual disabilities who share similar characteristics such as visual or cognitive impairment (PHE, 2019). However, there is a need for an increased understanding of the risk factors involved to enable the development of tailored interventions for this population group (Willgoss et al, 2010). A recent review by Pope et al (2021) aimed to build on previous findings, and explore the available evidence on risk factors for falls in people with ID. This commentary aims to critically appraise the methods used within the review by Pope et al (2021) and reflect on the applicability of these findings in community practice.

Methods

Pope et al (2021) conducted a narrative review to identify risk factors for falls among people with ID, building on previous findings of a similar review (Willgoss et al, 2010). A structured approach for documenting a search strategy was followed, incorporating key search terms related to falls, risk factors and ID. Literature searches were conducted using electronic databases (AMED, CINAHL, MEDLINE, PsychINFO) in November 2017 and again in July 2019. This was supplemented by searches of Google Scholar, the Cochrane Library and ASSIA. Studies included were those with: a population aged 18 years and over; adults with ID; reported outcomes on risk factors for falls; any geographical location or any accommodation type setting; participants living independently or with support; and those published in English in academic journals. Single case-studies, opinion pieces, commentaries, or those unable to describe a research methodology or published before February 2009, were excluded.

Screening of titles and full-text papers was undertaken by the first author and checked by the research team. Quality appraisal was undertaken by the research team using an author adapted scoring system of the Critical Appraisal Skills Programme (CASP), where a score of 17 or above represented strong quality and less than 10 reflected weak quality. A narrative synthesis was undertaken to draw out themes from both quantitative and qualitative data. Risk factors identified from more than one paper and including at least one significant association (reported Odds Ratio (OR) and a significance level of <0.05) were described. Thematic analysis was also undertaken to develop key themes from the data, including other risk factors identified (without a significant association reported), and checked by the research team.

Findings

After screening 705 records, eight studies were included in the review. Publication dates ranged from 2009 to 2019 and studies were conducted in Australia, Japan, Sweden,

Scotland, the Netherlands, New Zealand, and the US. Overall, 2506 participants were included in the review, of which, approximately 50% were male. A total of three studies included participants with mild to moderate ID, with none experiencing profound ID. Another study included a range from mild to profound ID, one study had participants with mostly severe to profound ID, and three studies did not specify the level of ID. Study settings varied from supported living to residential facilities, with no study conducted in a hospital setting. A total of five studies included quantitative data, one included qualitative data and a further two were mixed-methods studies. The review's assessment of bias for included studies rated the CASP scores between 13 to 18, which, according to the scoring system, indicated good qualities but with some methodological limitations. These included efforts made to minimise re-call bias, limited information on ethical approval, lack of detail as to how the studies' target samples were determined and how people with ID were involved in the data collection.

A total of nine risk factors were identified to have a significant association with falls in adults with ID. In addition, four main themes relating to potential risk factors for falls among people with ID were identified by the review: (1) the individual; (2) the situation; (3) factors for ongoing risks; and (4) protective factors that reduce falls (*Figure 1*).

The Individual

Risk factors that were found to have a significant association with falls and relating to the person, included: pre-existing conditions related to muscle weakness such as cerebral palsy (two quantitative papers); strength limitations as a result of underlying conditions such as arthritis and osteoporosis (one quantitative paper); history of previous falls (one quantitative paper); and epilepsy (three quantitative papers). Other potential factors identified in this theme included: increasing age; cognitive insight; and sex (females identified as being more at risk of falls).

The situation

Situational risk factors with a significant association included: unpredictability/impulsivity (one quantitative paper) and institutionalised environments compared to other types of accommodation (one quantitative paper). Other potential factors in this theme included: increased mobility; distractibility; location, for example, being indoors more so than outdoors and carers' vigilance (i.e., carers may be less protective if the person is indoors when vigilance is more relaxed); the home environment, due to taking more risks when encountering hazards such as people, furniture and temporary environmental changes; and being outdoors during autumn and winter.

Ongoing risks

Ongoing risk factors with a significant association included: urinary incontinence and rushing to the bathroom (two quantitative papers); engaging in vital activity of personal care such as toileting, eating or sleeping (one quantitative paper); and inappropriate or non-use of assistive equipment

(two quantitative papers). Other potential factors included: anticonvulsant medication (no reason provided, and a further study identified no differences between fallers and non-fallers for this factor); the role of carers in fall-management (consistency of support); and a recurring theme of the multifactorial nature of falls for people with ID, suggesting that falls do not occur due to isolated single risk factors.

Protective factors that reduce falls

Identified risk protectors included: carer vigilance and an awareness of environmental modifications to prevent falls, such as removing trip hazards and improving lighting; having Down's Syndrome (although no reasons given for this finding); people with Autism Spectrum Disorder (ASD) who may be less inclined to interact with the environment and in some instances, may be risk adverse; and people with moderate to severe ID with poor motor control who were more likely to be protected from falls by carers.

Commentary

Using the Joanna Briggs Institute's Critical Appraisal Tool for Systematic Reviews and Research Syntheses (Aromataris et al, 2015) to appraise the review conducted by Pope et al (2021), three of the 11 criteria were judged to be unclear. These criteria related to the use of critical appraisal and

data extraction. The authors utilised the CASP (2013) but failed to report which of the tools were used. An author adapted scoring system was also applied but this does not allow for weighting of individual questions and may therefore reflect an inaccurate portrayal of quality. It was also unclear if critical appraisal and data extraction were undertaken independently by the research team. For these reasons, the review findings, although comprehensive, should be interpreted with some caution when considering the implications for community practice.

The review's authors conclude that the number of risk factors identified from 2010 onwards has increased since Willgoss et al's (2010) review and that previously identified risk factors of increasing age, epilepsy, increased mobility and behavioural aspects were also recognised. New risk factors identified by Pope et al (2021) were related to the individual, the situation, and ongoing risk.

Working with this population in the community, an understanding of an individual's medical history could help to raise awareness of a heightened falls risk. The findings from this review suggests that the individual risk factors of decreasing physical ability, epilepsy, paretic conditions, and previous falls history should be taken note of within the patient's medical history due to their association with an increased risk of falls. It is also important to check that the




Figure 1: Potential risk factors and protectors.

individuals with ID have access to an annual health check as this can enhance their overall healthcare and reduce contributory risk factors for falls (Bakker-van Gijssel et al, 2017). As part of this appraisal process, it is important to ensure that individuals with ID also have access to a regular medication review and the option to take part in physical fitness and balance programme (Hale et al, 2016; O'Dwyer et al, 2018).

Further ongoing risk factors that should be identified when reviewing the individual's medical history are urinary incontinence and rushing to the bathroom; engaging in personal care, toileting, eating or sleeping; and inappropriate or non-use of assistive equipment. If these risk factors are identified, the use of an educational program on assisted aids, environmental adaptations such as a review of toilet access, strength programmes, and a review of personal care safety may be necessary to reduce the impact of these increased risks (Finlayson, 2018). Similarly, education on falls prevention for patients, family, and carers and/or a referral to a multi-disciplinary falls-prevention clinic may be warranted (Smulders et al, 2013; Finlayson, 2018).

These review findings also suggest that the situational risk factors of unpredictability/impulsivity, and institutionalised environments should be considered as an important factor associated with an increased risk of a fall. To counteract some of these risks, the use of home adaptations, mobility aids and assistive technology may be considered (Bainbridge et al, 2017; Finlayson, 2018).

Identification of fall risk factors in individuals with ID is essential due to the multifactorial and dynamic nature of falls (Ho et al, 2019). To help those in public health, health services and social care, Public Health England have produced guidance on preventing falls in people with learning disabilities (PHE, 2019). This tailored guidance suggests that possible contributors to falls should be considered in an individualised risk assessment, including internal factors such as health, vision, and fitness, behaviour and lifestyle, and external factors relating to environmental hazards, both at home and outside. Pro-active primary prevention is stated as useful, promoting a healthy lifestyle, including physical activity (with risk assessment) and regular health checks including vision, hearing and medication. More detailed suggestions for preventative strategies and risk assessments can be found in the guidance (PHE, 2019).

A multi-disciplinary team approach is important in preventing falls by people with ID (Finlayson, 2018). Community nurses may identify people with ID living in the community, who are at risk of falls. They can then work alongside other healthcare professionals such as occupational therapists (OTs), physiotherapists, general practitioners, and carers to provide tailored and multidisciplinary falls-prevention interventions. A referral by a member of the multi-disciplinary team, such as a community nurse, to an OT may be beneficial for assessment of individual and environmental risk factors. OTs can also tailor falls and injury prevention strategies for people with ID and monitor the use of assistive technology (Finlayson et al, 2014). This monitoring is important to ensure that aids and adaptations

are used safely and maintained regularly to prevent injury (PHE, 2019).

Recommendations

There is a need for further research to increase healthcare professionals' understanding of the range of individual, situational and ongoing risk factors for falls in people with ID, and subsequent falls-prevention interventions.

Community nurses should provide guidance on health-related issues associated with falls, and work in multidisciplinary healthcare teams as well as with carers of people with ID to provide falls-prevention interventions for individuals with ID at risk of falls.

Conclusions

Due to the limited amount of literature available on risk factors for falls among people with ID, and the wide variation in studies' samples, there is still a need to improve the research evidence on this topic. Age and sex were suggested fall risk factors for people with ID, and some protective factors for people with ID were also mooted by the available evidence, but further research is needed to confirm the limited studies' findings and the underlying reasons. Other areas for further research include addressing the need for standardised screening tools for people with ID who may have difficulty responding to self-assessment questions without support (Maring et al, 2017). Finally, there is a corresponding need to develop multi-component fall-prevention interventions for people with ID to address the wide variety of risk factors involved. Any future research should strive to include people with ID (and not just include those with mild-to-moderate ID), their families, carers, and supporters. **BJCN**

Acknowledgments: This commentary is an independent research partly funded by the National Institute for Health and Care Research (NIHR), Applied Research Collaboration North-West Coast (ARC NWC). The views expressed in this publication are those of the authors and not necessarily

Key points

- Falls are a serious and common problem for people with Intellectual Disability (ID)
- Risk factors for falls among people with ID are multifactorial and relate to the individual, situation, and ongoing risks
- Factors that protect against falls include carer vigilance and awareness in modifying the environment to remove traditional fall risks
- Further research is needed to develop falls prevention interventions for this population.

CPD reflective questions

- What are the differing risk factors for falls in people with ID compared to those in the general population and how could they be managed in the community?
- What interventions could be developed for addressing the multifactorial risk factors involved in falls among people with ID?

of those of the NIHR, the National Health Service, or the Department of Health and Social Care.

Accepted for publication: December 2022

Declaration of interest: None

Ang SGM, O'Brien AP, Wilson A. Carers' concern for older people falling at home: an integrative review. *Singapore Med J.* 2020;61(5):272-280. <https://doi.org/10.11622/smedj.2019142>

Aromataris E, Fernandez R, Godfrey C, Holly C, Kahlil H, Tungpukom P. Summarizing systematic reviews: methodological development, conduct and reporting of an Umbrella review approach. *Int J Evid Based Health.* 2015;13(3):132-140. <https://doi.org/10.1097/xeb.0000000000000055>

Bainbridge D, Hale L, Renfro M, Maring J. Addressing the Issue of Fall Risk for All Adults with Intellectual Developmental Disabilities: Part III: Creating Solutions with Community Fall Prevention and Inclusive Recreation Programs for Adults with Intellectual Developmental Disabilities. *GeriNotes.* 2017;24(5):16-20.

Bakker-van Gijssel EJ, Lucassen PLBJ, Olde Hartman TC et al. Health assessment instruments for people with intellectual disabilities-A systematic review. *Res Dev Disabil.* 2017;64:12-24. <https://doi.org/10.1016/j.ridd.2017.03.002>

Cox CR, Clemson L, Stancliffe RJ, Durvasula S, Sherrington C. Incidence of and risk factors for falls among adults with an intellectual disability. *J Intellect Disabil Res.* 2010;54(12):1045-1057. <https://doi.org/10.1111/j.1365-2788.2010.01333.x>

Critical Skills Appraisal Programme. CASP Checklists. 2013. <https://casp-uk.net/casp-tools-checklists/> (accessed 4 April 2023)

Finlayson J, Morrison J, Jackson A, Mantry D, Cooper S-A. Injuries, falls and accidents among adults with intellectual disabilities. Prospective cohort study. *J Intellect Disabil Res.* 2010;54(11):966-998. <https://doi.org/10.1111/j.1365-2788.2010.01319.x>

Finlayson J, Morrison J, Skelton DA, Ballinger C, Mantry D, Jackson A, Cooper S-A. The circumstances and impact of injuries on adults with learning disabilities. *The British Journal of Occupational Therapy.* 2014;77(8):400-409. <https://doi.org/10.4276/030802214X14071472109833>

Finlayson J. Injury and fall prevention for people with learning disabilities: a resource guide for people who care for, or support, people with learning disabilities. 2016. <https://agile.csp.org.uk/system/files/injury-and-fall-prevention-for-people-with-learning-disabilities-resource-guide.pdf> (accessed 4 April 2023)

Finlayson J. Fall prevention for people with learning disabilities: key points and recommendations for practitioners and researchers. *Tizard Learning Disability Review.* 2018; 23(2):91-99. <https://doi.org/10.1108/TLDR-06-2017-0026>

Hale LA, Mirfin-Veitch BF, Treharne GJ. Prevention of falls for adults with intellectual disability (PROFAID): a feasibility study. *Disabil Rehabil.* 2016;38(1):36-44. <https://doi.org/10.3109/09638288.2015.1017613>

Hermans H, Evenhuis HM. Multi-morbidity in older adults with intellectual disabilities. *Res Dev Disabil.* 2014;35(4):776-83. <https://doi.org/10.1016/j.ridd.2014.01.022>

Ho P, Bulsara M, Patman S, Downs J, Bulsara C, Hill AM. Incidence and associated risk factors for falls in adults with intellectual disability. *J Intellect Disabil Res.* 2019;63(12):1441-1452. <https://doi.org/10.1111/jir.12686>

Hsieh K, Rimmer J, Heller T. Prevalence of falls and risk factors in adults with intellectual disability. *Am J Intellect Dev Disabil.* 2012;117:442-454. <https://doi.org/10.1352/1944-7558-117.6.442>

Maring J, Hale L, Renfro M, Bainbridge D. Addressing the issue of fall risk for all adults with intellectual developmental disabilities. Part II. *Geri Notes.* 2017;24(5):12-15.

Mencap. How common is learning disability in the UK? <https://www.mencap.org.uk/learning-disability-explained/research-and-statistics/how-common-learning-disability/> (accessed 4 April 2023)

National Institute for Health and Care Excellence. Falls in older people: assessing risk and prevention. Clinical guideline. 2013. <https://www.nice.org.uk/guidance/cg161> (accessed 4 April 2023)

O'Dwyer M, McCallion P, McCarron M, Henman M. Medication use and potentially inappropriate prescribing in older adults with intellectual disabilities: a neglected area of research. *Ther Adv Drug Saf.* 2018;9(9):535-557. <https://doi.org/10.1177/2042098618782785>

Oppewal A, Hilgenkamp TI, van Wijck R, Schoufour JD, Evenhuis HM. The predictive value of physical fitness for falls in older adults with intellectual disabilities. *Res Dev Disabil.* 2014;35(6):1317-1325. <https://doi.org/10.1016/j.ridd.2014.03.022>

Pal J, Hale L, Mirfin-Veitch B, Claydon L. Injuries and falls among adults with intellectual disability: a prospective New Zealand cohort study. *J Intellect Disabil Res.* 2013;39:35-44. <https://doi.org/10.3109/13668250.2013.867929>

Petropoulou, E, Finlayson, J, Hay, M, Spencer, W, Park, R, Tannock, H, Galbraith, E, Godwin, J, & Skelton, D.A. Injuries reported and recorded for adults with intellectual disabilities who live with paid support in Scotland: a comparison with Scottish adults in the general population. *J Appl Res Intellect Disabil.* 2017;30(2):408-415. <https://doi.org/10.1111/jar.12244>

Pope, J, Truesdale, M, Brown, M. Risk factors for falls among adults with intellectual disabilities: A narrative review. *J Appl Res Intellect Disabil.* 2021;34:274-285. <https://doi.org/10.1111/jar.12805>

Public Health England (PHE). 2019. Preventing falls in people with learning disabilities: making reasonable adjustments. <https://www.gov.uk/government/publications/preventing-falls-in-people-with-learning-disabilities/preventing-falls-in-people-with-learning-disabilities-making-reasonable-adjustments#fn:2> (accessed 4 April 2023)

Smulders E, Enkelar L, Schoon Y, Geurts AC, van Schroyen Lantman-de Valk H, Weerdesteijn V. Falls prevention in persons with intellectual disabilities: development, implementation, and process evaluation of a tailored multifactorial fall risk assessment and intervention strategy. *Res Dev Disabil.* 2013;34(9):2788-2798. <https://doi.org/10.1016/j.ridd.2013.05.041>

Snooks H, Cheung WY, Gwini SM, Humphreys I, Sanchez A, Sirwardena N. Can older people who fall be identified in the ambulance call centre to enable alternative responses or care pathways? *Emerg Med J.* 2011;28(3)e1. <http://dx.doi.org/10.1136/emj.2010.108605.9v>

World Health Organization. Better health, better lives: children and young people with intellectual disabilities and their families. 2010. https://www.euro.who.int/__data/assets/pdf_file/0003/125751/e94414.pdf (accessed 4 April 2023)

World Health Organization. Factsheet: falls. 2021. <https://www.who.int/news-room/fact-sheets/detail/falls> (accessed 4 April 2023)

Willgoss TG, Yohannes AM, Mitchell D. Review of risk factors and preventative strategies for fall related injuries in people with intellectual disabilities. *J Clin Nurs.* 2010;19(15-16):2100-2109. <https://pubmed.ncbi.nlm.nih.gov/20569280/>

A-Z Dictionary of Wound Care

Fiona Collins, Sylvie Hampton, Richard White

- Essential dictionary defining words and terms that are used in the field of tissue viability
- Essential guide for students or those aspiring to become specialists
- Includes more rarely used terms

ISBN: 978-1-85642-225-3; 216 x 138 mm; paperback; 112 pages; publication 2002; £19.99

Order your copies by visiting
www.quaybooks.co.uk

or call our Hotline
+44(0)1722 716935

